

DEMOGRAPHICS

Patient Name: _____
Address: _____
City: _____ State: _____ Zip: _____
DOB: ____/____/____ ☐ Male ☐ Female
Phone: _____
SSN: _____ Ht: _____ Wt: _____

PRESCRIBING PHYSICIAN

Name: _____
Address: _____
City: _____ State: _____ Zip: _____
Phone: _____ Fax: _____
License#: _____
DEA#: _____ NPI: _____

ALLERGIES / REACTIONS

Signature: _____
Date: _____

PHYSICIAN ORDERS

☐ **BENLYSTA (BELIMUMAB)**

REFILLS: _____

☐ Infuse 10 mg/kg in 250 ml of NS IV over 60 minutes

FREQUENCY: ☐ Induction: 0, 14 days, 28 days ☐ Every 28 days

☐ **IV ACCESS**

☐ Peripheral IV - Flush with Normal Saline 5–10 ml with IV start and before & after medication

☐ Subcutaneous Port - Flush with Normal Saline 10 ml before & after medication—LOCK with Heparin 500 units (100 units/ml)

☐ **PRE-MEDICATIONS / LABS**

REFILLS: _____

☐ Diphenhydramine ____ MG IV 30 minutes before infusion ☐ Solu-Medrol ____ MG IV 30 minutes before infusion

☐ OTHER: _____

☐ LABS: _____ FREQUENCY _____

☒ **Anaphylaxis Kit per Pharmacy protocol**

TO BE ADMINISTERED BY THE HEALTH CARE PROVIDER PRN FOR PATIENTS RECEIVING IV MEDS IN THE HOME
Epinephrine Auto Injector 0.3mg/0.3ml IM – Repeat one time in 20 minutes if needed

☒ **ADVERSE REACTION ORDERS:**

Mild Reactions - Reduce rate by 1/2 at time of onset and keep reduced x30 minutes. If resolved, increase per guidelines. If not resolved, administer appropriate medication based on symptoms,

Severe Reactions - STOP infusion ☒, KVO IV Normal Saline ☒ Diphenhydramine 25/50 mg IVP over 3-5 minutes ☒
Methylprednisolone 125 mg IVP over 5 minutes ☒ Call Ambulance and Physician ☒.

DIAGNOSIS:

☐ M32.9 Moderate to Severe Systemic Lupus Erematosus (SLE)

☐ Other: _____