

DEMOGRAPHICS

Patient Name: _____ Name: _____
Address: _____ Address: _____
City: _____ State: _____ Zip: _____ City: _____ State: _____ Zip: _____
DOB: ____/____/____ ☐ Male ☐ Female Phone: _____ Fax: _____
Phone: _____ License#: _____
SSN: _____ Ht: _____ Wt: _____ DEA#: _____ NPI: _____

PRESCRIBING PHYSICIAN

Signature: _____
Date: _____

ALLERGIES / REACTIONS

PHYSICIAN ORDERS

☐ **IMMUNOGLUBLIN – IV** _____ **REFILLS:** _____

INFUSE _____ G/KG over _____ days – REPEAT every _____ WEEKS

- ☐ Peripheral IV - Flush with Normal Saline 5–10 ml with IV start and before & after medication
☐ Subcutaneous Port - Flush with Normal Saline 10 ml before & after medication.
LOCK with Heparin 500 units (100 units/ml)

☐ **PRE-MEDICATIONS / LABS** **REFILLS:** _____

- ☐ Diphenhydramine 25 MG IV 30 minutes before infusion
☐ Solu-Medrol 125 MG IV – 30 minutes before infusion
☐ OTHER: _____
☐ LABS: _____

☒ **Anaphylaxis Kit per Pharmacy protocol**

TO BE ADMINISTERED BY THE HEALTH CARE PROVIDER **PRN** FOR PATIENTS RECEIVING IV MEDS IN THE HOME
Epinephrine Auto Injector 0.3mg/0.3ml IM – Repeat one time in 20 minutes if needed

☒ **ADVERSE REACTION ORDERS:**

Mild Reactions - Reduce rate by 1/2 at time of onset and keep reduced x30 minutes. If resolved, increase per guidelines. If not resolved, administer appropriate medication based on symptoms,
Severe Reactions - STOP infusion, KVO IV Normal Saline _____ Diphenhydramine 25/50 mg IVP over 3-5 minutes _____
Methylprednisolone 125 mg IVP over 5 minutes _____ Call Ambulance and Physician.

DIAGNOSIS:

☐ _____ ☐ _____