

DEMOGRAPHICS

Patient Name: _____

Address: _____

City: _____ State: _____ Zip: _____

DOB: ____/____/____ ☐ Male ☐ Female

Phone: _____

SSN: _____ Ht: _____ Wt: _____

PRESCRIBING PHYSICIAN

Name: _____

Address: _____

City: _____ State: _____ Zip: _____

Phone: _____ Fax: _____

License#: _____

DEA#: _____ NPI: _____

ALLERGIES / REACTIONS

Signature: _____

Date: _____

PHYSICIAN ORDERS

☐

SIMPONI ARIA

REFILLS: _____

☐ Initial Dose: 2mg/kg (_____mg) IV at weeks 0 & 4

☐ Maintenance Dose: 2 mg/kg (_____mg) every 8 weeks

☐

IV ACCESS

☐

Peripheral IV - Flush with Normal Saline 5-10 ml with IV start and before & after medication

☐

Subcutaneous Port - Flush with Normal Saline 10 ml before & after medication---LOCK with Heparin 500 units (100 units/ml)

☐

PRE-MEDICATIONS / LABS

REFILLS: _____

☐

Diphenhydramine _____ MG IV 30 minutes before infusion

☐

Solu-Medrol _____ MG IV 30 minutes before infusion

☐

OTHER: _____

☐

LABS: _____ FREQUENCY _____

☒

Anaphylaxis Kit per Pharmacy protocol

TO BE ADMINISTERED BY THE HEALTH CARE PROVIDER PRN FOR PATIENTS RECEIVING IV MEDS IN THE HOME

Epinephrine Auto Injector 0.3mg/0.3ml IM - Repeat one time in 20 minutes if needed

☒

ADVERSE REACTION ORDERS:

Mild Reactions - Reduce rate by 1/2 at time of onset and keep reduced x30 minutes. If resolved, increase per guidelines. If not resolved, administer appropriate medication based on symptoms,

Severe Reactions - STOP infusion ☒, KVO IV Normal Saline ☒ Diphenhydramine 25/50 mg IVP over 3-5 minutes ☒

Methylprednisolone 125 mg IVP over 5 minutes ☒ Call Ambulance and Physician ☒.

DIAGNOSIS:

☐

M06.9 Rheumatoid Arthritis

☐

M08.0 Juvenile Rheumatoid Arthritis

☐

L40.50 Psoriatic Arthritis

☐

M45.9 Ankylosing Spondylitis

☐

Other: _____