

DEMOGRAPHICS

Patient Name: _____

Address: _____

City: _____ State: _____ Zip: _____

DOB: ____/____/____ ☐ Male ☐ Female

Phone: _____

SSN: _____ Ht: _____ Wt: _____

PRESCRIBING PHYSICIAN

Name: _____

Address: _____

City: _____ State: _____ Zip: _____

Phone: _____ Fax: _____

License#: _____

DEA#: _____ NPI: _____

ALLERGIES / REACTIONS

Signature: _____

Date: _____

PHYSICIAN ORDERS

☐

STELARA

REFILLS: _____

☐ 130 mg vial

☐ < 121 lbs — Infuse IV 260 mg (2 vials) over at least 1 hour

☐ > 121 lbs to 187 lbs — Infuse IV 390 mg (3 vials) over at least 1 hour

☐ > 188 lbs to — Infuse IV 520 mg (4 vials) over at least 1 hour

☐ 90 mg vial/syringe

☐ Inject 90 mg SQ 8 weeks after initial IV infusion then every 8 weeks thereafter

☐

IV ACCESS

☐ Peripheral IV - Flush with Normal Saline 5–10 ml with IV start and before & after medication

☐ Subcutaneous Port - Flush with Normal Saline 10 ml before & after medication—LOCK with Heparin 500 units (100 units/ml)

☐

PRE-MEDICATIONS / LABS

REFILLS: _____

☐ Diphenhydramine ____ MG IV 30 minutes before infusion ☐ Solu-Medrol ____ MG IV 30 minutes before infusion

☐ OTHER: _____

☐ LABS: _____ FREQUENCY _____

☒

Anaphylaxis Kit per Pharmacy protocol

TO BE ADMINISTERED BY THE HEALTH CARE PROVIDER PRN FOR PATIENTS RECEIVING IV MEDS IN THE HOME

Epinephrine Auto Injector 0.3mg/0.3ml IM – Repeat one time in 20 minutes if needed

☒

ADVERSE REACTION ORDERS:

Mild Reactions - Reduce rate by 1/2 at time of onset and keep reduced x30 minutes. If resolved, increase per guidelines. If not resolved, administer appropriate medication based on symptoms,

Severe Reactions - STOP infusion ☒, KVO IV Normal Saline ☒ Diphenhydramine 25/50 mg IVP over 3-5 minutes ☒

Methylprednisolone 125 mg IVP over 5 minutes ☒ Call Ambulance and Physician ☒.

DIAGNOSIS:

☐

K50.90 Crohn's Disease NOS

☐

K51.90 Ulcerative Colitis

☐

Other: _____