

DEMOGRAPHICS

Patient Name: _____
Address: _____
City: _____ State: _____ Zip: _____
DOB: ____/____/____ ☐ Male ☐ Female
Phone: _____
SSN: _____ Ht: _____ Wt: _____

PRESCRIBING PHYSICIAN

Name: _____
Address: _____
City: _____ State: _____ Zip: _____
Phone: _____ Fax: _____
License#: _____
DEA#: _____ NPI: _____

ALLERGIES / REACTIONS

Signature: _____

Date: _____

PHYSICIAN ORDERS



VENOFER 200 mg IV – 5 infusions over 14 days

REFILLS: _____

Infuse VENOFER 200 mg/0.9% NaCl 100 ml IV over 20-30 minutes



IV ACCESS



Peripheral IV - Flush with Normal Saline 5–10 ml with IV start and before & after medication



Subcutaneous Port - Flush with Normal Saline 10 ml before & after medication—LOCK with Heparin 500 units (100 units/ml)



PRE-MEDICATIONS / LABS

REFILLS: _____



Diphenhydramine ____ MG IV 30 minutes before infusion



Solu-Medrol ____ MG IV 30 minutes before infusion



OTHER: _____



LABS: _____ FREQUENCY _____



Anaphylaxis Kit per Pharmacy protocol

TO BE ADMINISTERED BY THE HEALTH CARE PROVIDER PRN FOR PATIENTS RECEIVING IV MEDS IN THE HOME

Epinephrine Auto Injector 0.3mg/0.3ml IM – Repeat one time in 20 minutes if needed



ADVERSE REACTION ORDERS:

Mild Reactions - Reduce rate by 1/2 at time of onset and keep reduced x30 minutes. If resolved, increase per guidelines. If not resolved, administer appropriate medication based on symptoms,

Severe Reactions - STOP infusion ☒, KVO IV Normal Saline ☒ Diphenhydramine 25/50 mg IVP over 3-5 minutes ☒

Methylprednisolone 125 mg IVP over 5 minutes ☒ Call Ambulance and Physician ☒.

DIAGNOSIS:



D50.9 Iron deficiency anemia, unspecified



E61.1 Iron deficiency