

DEMOGRAPHICS

Patient Name: _____

Address: _____

City: _____ State: _____ Zip: _____

DOB: ____/____/____ ☐ Male ☐ Female

Phone: _____

SSN: _____ Ht: _____ Wt: _____

PRESCRIBING PHYSICIAN

Name: _____

Address: _____

City: _____ State: _____ Zip: _____

Phone: _____ Fax: _____

License#: _____

DEA#: _____ NPI: _____

ALLERGIES / REACTIONS

Signature: _____

Date: _____

PHYSICIAN ORDERS

☐

SPEVIGO

REFILLS: _____

☐ **900 mg IV over 90 minutes** - If flare symptoms persist, an additional 900 mg dose of Spevigo may be administered one week after the initial dose. If needed, please submit a separate order form for this dose

☐ **600 mg SQ, then 300 mg SQ** every 4 weeks

☐ **300 mg SQ every 4 weeks** (for patients that have received 900 mg IV dose 4 weeks prior)

☐

IV ACCESS

☐

Peripheral IV - Flush with Normal Saline 5-10 ml with IV start and before & after medication

☐

Subcutaneous Port - Flush with Normal Saline 10 ml before & after medication---LOCK with Heparin 500 units (100 units/ml)

☐

PRE-MEDICATIONS / LABS

REFILLS: _____

☐

Diphenhydramine ____ MG IV 30 minutes before infusion

☐

Solu-Medrol ____ MG IV 30 minutes before infusion

☐

OTHER: _____

☐

LABS: _____ FREQUENCY _____

☒

Anaphylaxis Kit per Pharmacy protocol

TO BE ADMINISTERED BY THE HEALTH CARE PROVIDER PRN FOR PATIENTS RECEIVING IV MEDS IN THE HOME

Epinephrine Auto Injector 0.3mg/0.3ml IM - Repeat one time in 20 minutes if needed

☒

ADVERSE REACTION ORDERS:

Mild Reactions - Reduce rate by 1/2 at time of onset and keep reduced x30 minutes. If resolved, increase per guidelines. If not resolved, administer appropriate medication based on symptoms,

Severe Reactions - STOP infusion ☒, KVO IV Normal Saline ☒ Diphenhydramine 25/50 mg IVP over 3-5 minutes ☒

Methylprednisolone 125 mg IVP over 5 minutes ☒ Call Ambulance and Physician ☒.

DIAGNOSIS:

☐

☐
